

## ROLE OF HYPERBILIRUBINEMIA : A NEW DIAGNOSTIC TOOL ANDA PREDICTOR OF GANGRENOUS / PERFORATED APPENDICITIS IN PAEDIATRIC POPULATION

Udhayasankar V<sup>1</sup>, Haris Vijila Rani M<sup>2</sup>

<sup>1</sup>Assistant Professor, Department of Paediatric Surgery, Government Vellore Medical College Hospital, Vellore, Tamil Nadu, India

<sup>2</sup>Associate Professor, Department of Paediatric Surgery, Government Vellore Medical College Hospital, Vellore, Tamil Nadu, India

Received : 19/12/2023  
Received in revised form : 21/02/2024  
Accepted : 04/03/2024

**Keywords:**

Serum Bilirubin, Acute Appendicitis, Acute Complicated Appendicitis.

**Corresponding Author:**

Dr. Udhayasankar V,  
Email: drudhayasankar07@gmail.com

DOI: 10.47009/jamp.2024.6.2.138

Source of Support: Nil,  
Conflict of Interest: None declared

Int J Acad Med Pharm  
2024; 6 (2); 652-654



### Abstract

**Background:** Acute Appendicitis is one of the most common surgical emergency encountered in day to day emergency surgical practice. Delay in diagnosis and treatment can lead to gangrene, perforation and diffuse peritonitis and acute/subacute intestinal obstruction in paediatric population. The aim and objective is to find out the specificity, sensitivity, predictive value of positive test and predictive value of negative test of Serum Bilirubin in diagnosis of acute appendicitis and acute complicated appendicitis. **Materials and Methods:** All the patients admitted with clinical diagnosis of Acute Appendicitis were tested by laboratory investigations and ultrasonography of the abdomen. Preoperatively patient's blood was also collected for serum bilirubin and other liver enzymes estimation. Cases that underwent emergency appendectomy from September 2021 - August 2023 were included in present Prospective Analytical Study. **Result:** About 100 patients were included in this prospective analytical study. Hyperbilirubinemia as a sensitivity of 84.7% and specificity of 80%, Predictive Value of positive test is 96%, Predictive value of negative test is 48%. In Complicated appendicitis that is gangrenous/perforated Appendicitis, the serum bilirubin levels higher compared to the acute appendicitis. **Conclusion:** The level of serum bilirubin aids in the diagnosis of Acute Appendicitis and higher levels as a predictive potential for the Acute Complicated Appendicitis(Gangrenous/Perforated).

## INTRODUCTION

Acute appendicitis is the most common surgical emergency seen in day-to-day practice in emergency department.<sup>[1]</sup> It can sometimes confuse the practitioners by its presentation. The delay in early diagnosis or failure in early diagnosis may happen many times.<sup>[2]</sup> This may lead on to the poor disease prognosis. This will further lead on to increase in morbidity as well as occasional mortality in the patient though there are many recent trends in investigatory modalities.

Diagnosis of acute appendicitis is still a mystery. There is increase in the negative appendectomy rate of about 15 - 30 % seen in literature.<sup>[3]</sup>

So number of various investigations were used to reduce the rate of negative appendectomy such as Laboratory investigations and Radiological investigations.<sup>[4]</sup>

### Objectives

To find out the specificity, sensitivity, predictive value of positive test and predictive value of negative

test of Serum Bilirubin in diagnosis of acute appendicitis and acute complicated appendicitis.

## MATERIALS AND METHODS

This study conducted as Prospective Analytical Study.

This study was conducted in 100 patients. These 100 patients are clinically diagnosed to have acute appendicitis, and who were underwent emergency appendectomy. This study conducted in Govt.Vellore Medical college Hospital,Vellore, Department of Paediatric surgery, during the period from September 2021 to August 2023.

Serum Bilirubin of more than 1.6mg/dl considered high and more than 2.2mg/dl considered very high.

### Inclusion Criteria

All the patients below 18 years of age diagnosed to have appendicitis included in this study and subjected for appendectomy in Govt.Vellore Medical college, Vellore.

### Exclusion Criteria

Patients who were HBsAg positive, patients taking hepatotoxic drugs, where serum bilirubin was raised

Sensitivity =  $TP \times 100 / TP + FN$

Specificity =  $TN \times 100 / TN + FP$

Predictive Value of Positive Test =  $TP \times 100 / TP + FP$

Predictive Value Of Negative Test =  $TN \times 100 / TN + FN$

## RESULTS

In 100 patients were included in this study and underwent Emergency Appendicectomy.

Age distribution

The acute appendicitis fall in to 5 groups in the study population. In this study maximum incidence is seen in 3rd group of life that is the age group between 10 to 14 years, followed by the age group between 6 to 49 yrs, least incidence of appendicitis occurrence seen in infants age group shown in [Table 1].




Figure 1: Serum Bilirubin with HPE Correlation

### Sex Distribution

In our study the number of male children are 55, and female children are 45 in number.

[Table 2] depicts, In present study out of 100 patients 85 patients has HPE positive, among 85 positive individuals serum bilirubin was raised in 72 patients that is 84.7%. 13 patients had normal serum

bilirubin value ( $x^2$  Mc nemor = 8.9,  $P = < 0.01$ ) significant association in diagnosis of acute appendicitis.

In this study among 14 complicated patients, 13 patients had very high raise in serum bilirubin value 92%, where as only 1 patients had very high serum bilirubin value in uncomplicated appendicitis.

In present study serum bilirubin shows specificity of 80%, predictive value of positive test 96%, Negative predictive value 48%.




Figure 2: Distribution of cases according to Serum Bilirubin Value.




Figure 3: Sensitivity, Specificity, Predictive Value of Positive Test and Negative Test of Serum Bilirubin

Table 1: Age Distribution

Age(years)	Incidence(outof 100)	Percentage
Infants 0 -1 year	1	1%
1-5 years	15	15%
6 – 9 years	28	28%
10 -14 years	32	32%
>14yrs – <18 years	24	24%
Total	100	100%

Table 2:Role Of Serum Bilirubin

Serum bilirubin	HPE		Total
	Positive	Negative	
Positive	72TP	3FP	75
Negative	13FN	12TN	25
Total	85	15	100

## DISCUSSION

In this study it reveals that serum bilirubin is highly sensitive in diagnosis of complicated appendicitis sensitivity of 92%. The sensitivity of serum bilirubin increase in uncomplicated appendicitis is 84.7%.

The interesting finding of this study was that the positive predictive value was 96%, which aids in the diagnosis of acute appendicitis, if the serum bilirubin goes beyond >2.2mg/dl, the possibility of complicated appendicitis is very high.

Bilirubin is not commonly known to be a relevant marker in appendicitis. However, previous studies have found hyperbilirubinaemia to be a marker with high specificity for perforated appendicitis. Adult bilirubin levels in the adult surgical population are usually raised due to liver or gallbladder problems.

Hepatic dysfunction as a result of bacterial infection or sepsis without direct invasion of the liver, has already been well described. The possible aetiologies include gram negative sepsis, shock and ischaemic liver, severe trauma or gut barrier failure. Gram negative organisms, with *Escherichia coli* in particular, have been shown to produce endotoxins that affect bile flow in rat liver models. None of the patients in our study had ischaemic liver, or septic shock.<sup>[6,7]</sup>

In appendicitis, compromised appendix wall integrity leads to translocation of bacteria and endotoxins from the appendix lumen into the portal system. Inflammatory cytokines may then travel to the liver, inducing intrahepatic cholestasis. Research has also shown that *E. coli* endotoxin causes dose dependent cholestasis, which would explain our findings of increased bilirubin levels with progressive appendicitis severity. It is possible that bilirubin may be raised in other sources of gram-negative related sepsis which may be of a gastrointestinal origin (*E. coli* in diverticulitis) or from other sources such as pneumonia, endocarditis, pyelonephritis and soft-tissue abscesses.<sup>[8]</sup>

Estrada et al,<sup>[9]</sup> 2007 in his Retrospective study of 170 patients with findings of Hyperbilirubinaemia (17.1 mmol/L) and perforated appendicitis  $p < 0.031$ , Odds ratio 2.96 (95% confidence interval 1.11e7.6)

Sand et al,<sup>[10]</sup> 2009 in his Retrospective study of 538 patients with Hyperbilirubinaemia (26.5 mmol/L) and perforated appendicitis  $p < 0.05$ , Specificity 86%

Kaser et al,<sup>[11]</sup> 2010 in his Retrospective study of 725 patients Hyperbilirubinaemia (20 mmol/L) and perforated appendicitis  $p < 0.05$ , Specificity 78%

Atahan et al,<sup>[12]</sup> 2011 in his Retrospective study of 351 patients Hyperbilirubinaemia (17.1 mmol/L) and perforated appendicitis  $p = 0.000$ , Specificity 87.21%.

Emmanuel et al,<sup>[13]</sup> 2011 in his Retrospective study of 472 patients Hyperbilirubinaemia (20.5 mmol/L)

and simple appendicitis  $p < 0.001$ , Specificity 88% Hyperbilirubinaemia and perforated appendicitis  $p < 0.001$ , Specificity 70%.

Beltran et al,<sup>[14]</sup> 2009 in his Prospective study of 134 patients Hyperbilirubinaemia (17.86 mmol/L) and perforated appendicitis Specificity 51%, Sensitivity 57%

Khan et al,<sup>[15]</sup> 2004 observed among 110 individuals, the serum bilirubin value was increased in acute appendicitis, it is more specifically elevated in complicated appendicitis. Specificity of this study in complicated appendicitis is 80.6%, in uncomplicated acute appendicitis 56.36%.

## CONCLUSION

Bilirubin is a specific marker for acute appendicitis with a good positive predictive value. It is also a valuable indicator of patients more likely to have appendiceal perforation or gangrene. Bilirubin should be used together with clinical examination and other laboratory investigations in the assessment of patients with suspected acute appendicitis.

## REFERENCES

1. Peranteau WH, Smink DS. Appendix, Meckel's and other small bowel diverticula. In: Zinner MJ, Ashley W. Stanley, editors. *Maingot's Abdominal Operation*. 12th ed. New York: The McGraw-Hill Companies; 2013. p. 623-40.
2. Alvarado A.: A practical score for the early diagnosis of acute appendicitis. *Ann Emerg Med* 1986; 15: 557-564
3. Chang F.C., Hogle H.H., Welling D.R.: The fate of the negative appendix. *Am. J. Surg.* 1973;126:752-754
4. Pearson R.H. Ultrasonography for diagnosing appendicitis. *Br Med. J.* 1988;297:309-310
5. Chand N, Sanyal AJ. Sepsis-induced cholestasis. *Hepatology* (Baltimore, Md) 2007 Jan;45(1):230e41. PubMed PMID: 17187426. Epub 2006/12/26. eng.
6. Utili R, Abernathy C, Zimmerman H. Studies on the effects of *C. coli* endotoxin on canalicular bile formation in the isolated perfused rat liver. *J Lab Clin Med* 1977;89(3):471.
7. Utili R, Abernathy CO, Zimmerman H. Inhibition of Na<sup>+</sup>, K<sup>+</sup>-adenosine-triphosphatase by endotoxin: a possible mechanism for endotoxin-induced cholestasis. *J Infect Dis* 1977;136(4):583e7.
8. Rauner M, Fickert P, Stauber RE. Inflammation-induced cholestasis. *J Gastro-enterol Hepatol* 1999;14(10):946e59.
9. Estrada JJ, Petrosyan M, Barnhart J, Tao M, Sohn H, Towfigh S, et al. Hyper-bilirubinemia in appendicitis: a new predictor of perforation. *J Gastrointest Surg* 2007 Jun;11(6):714e8. PubMed PMID: 17436050. Epub 2007/04/17. eng.
10. Sand M, Bechara FG, Holland-Letz T, Sand D, Mehnert G, Mann B. Diagnostic value of hyperbilirubinemia as a predictive factor for appendiceal perforation in acute appendicitis. *Am J Emerg Med* 2009 Aug;19(2):193e8. PubMed PMID: 19306980. Epub 2009/03/25. eng.
11. Kaser SA, Fankhauser G, Willi N. C.A.Maurer. C-reactive protein is superior to bilirubin for anticipation of perforation in acute appendicitis. *Scand J Gastro-enterol* 2010 Aug;45(7-8):885e92.
12. Atahan K, Ureyen O, Aslan E, Deniz M, Cokmez A, Gur S, et al. Preoperative diagnostic role of hyperbilirubinaemia as a marker of appendix perforation. *J Int Med Res.* 2011;39(2):609e18. PubMed PMID: 21672366. Epub 2011/06/16. eng.
13. Emmanuel A, Murchan P, Wilson I, Balfe P. The value of hyperbilirubinaemia in the diagnosis of acute appendicitis. *Ann R Coll Surg Engl* 2011;93(3):213e7.
14. Beltran M, Mendez P, Barrera R, Contreras M, Wilson C, Cortes V, et al. Is hyperbilirubinaemia in appendicitis a better predictor of perforation than C-reactive protein? A prospective study. *Indian J Surg* 2009;71(5):265e72.
15. Khan S. Elevated serum bilirubin in acute appendicitis: a new diagnostic tool. *Kathmandu Univ Med J (KUMJ)* 2008 Apr-Jun;6(2):161e5. PubMed PMID: 18769079. Epub 2008/09/05. eng.